

ON A LOWER BOUND OF THE KOBAYASHI METRIC

NIKOLAI NIKOLOV

ABSTRACT. It is shown that a lower bound of the Kobayashi metric of convex domains in $\mathbb{C}^n$ does not hold for non-convex domains.

Let D be a domain in $\mathbb{C}^n$, $z \in D$ and $X \in \mathbb{C}^n$. Denote by γ_D and κ_D the Carathéodory and Kobayashi metrics of D :

$$\gamma_D(z; X) = \sup\{|f'(z)X| : f \in \mathcal{O}(D, \mathbb{D})\},$$

$$\kappa_D(z; X) = \inf\{|\alpha| : \exists \varphi \in \mathcal{O}(\mathbb{D}, D) \text{ with } \varphi(0) = z, \alpha\varphi'(0) = X\},$$

where $\mathbb{D}$ is the unit disc. Then $\gamma_D \leq \kappa_D$. By the Lempert theorem [5], $\gamma_D = \kappa_D$ if D is convex.

Set $d_D(z; X)$ to be the distance from z to ∂D in the X -direction, i.e.

$$d_D(z; X) = \sup\{r > 0 : z + \lambda X \in D \text{ if } |\lambda| < r\}$$

(possibly $d_D(z; X) = \infty$). It follows by the definition of κ_D that

$$\kappa_D(z; X) \leq \frac{1}{d_D(z; X)}.$$

On the other hand, if D is convex, then

$$\gamma_D(z; X) \geq \frac{1}{2d_D(z; X)}$$

due to S. Frankel [2, Theorem 2.2] and I. Graham [3, Theorem 5]. The following short proof of this estimate can be found in [1, Theorem 4.1]:

$$\gamma_D(z; X) \geq \gamma_\Pi(z; X) = \frac{1}{2d_\Pi(z; X)} = \frac{1}{2d_D(z; X)},$$

where Π is a supporting half-space of D at a boundary point of the form $z + \lambda X$, $|\lambda| = d_D(z, X)$.

It turns out that the converse to the Frankel-Graham result holds.

Proposition 1. *For a domain D in $\mathbb{C}^n$ the following conditions are equivalent:*

(a) D is convex;

(b) $\gamma_D(z; X) \geq \frac{1}{2d_D(z; X)}$;

2010 *Mathematics Subject Classification.* 32F45.

Key words and phrases. Carathéodory and Kobayashi metrics, convex domain.

$$(c) \liminf_{z \rightarrow a} \frac{2\kappa_D(z; z - a) - 1}{\|z - a\|} \geq 0 \text{ for any } a \in \partial D.$$

Proof. We have only to show that (c) implies (a). Assume the contrary. According to [4, Theorem 2.1.27], one may find a point $a \in \partial D$ and a real-valued quadratic polynomial q such that $q(a) = 0$, $\nabla q(a) \neq 0$, the set $G = \{z \in \mathbb{C}^n : q(z) < 0\}$ is contained in D near a and ∂G has normal curvature $\chi < 0$ in some direction $X \in T_a^{\mathbb{R}} \partial G$. We may assume by continuity that $X \notin T_a^{\mathbb{C}} \partial G$. Then the planar set $F = G \cap (a + \mathbb{C}X)$ has smooth boundary near a with curvature χ at a . Let E be the connected component of F for which $a \in \partial E$. Denote by n the inner normal to ∂E at a . Using [6, Proposition 1], we get the contradiction

$$0 \leq \limsup_{n \ni z \rightarrow a} \frac{2\kappa_D(z; z - a) - 1}{\|z - a\|} \leq \lim_{n \ni z \rightarrow a} \frac{2\kappa_E(z; z - a) - 1}{\|z - a\|} = \frac{\chi}{2} < 0.$$

REFERENCES

- [1] E. Bedford, S. I. Pinchuk, *Convex domains with noncompact groups of automorphisms*, Mat. Sb. 185(5) (1994), 3-26.
- [2] S. Frankel, *Applications of affine geometry to geometric function theory in several complex variables Part I. Convergent rescalings and intrinsic quasi-isometric structure*, Proc. Symp. Pure Math. 52(2) (1991), 183-208.
- [3] I. Graham, *Sharp constants for the Koebe theorem and for estimates of intrinsic metrics on convex domains*, Proc. Symp. Pure Math. 52(2) (1991), 233-238.
- [4] L. Hörmander, *Notions of convexity*, Birkhäuser, Boston, 1994.
- [5] L. Lempert, *Holomorphic retracts and intrinsic metrics in convex domains*, Anal. Math. 8 (1982), 257-264.
- [6] N. Nikolov, M. Trybula, L. Andreev, *Boundary behavior of invariant functions on planar domains*, arXiv:1511.03117.

INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, ACAD. G. BONCHEV 8, 1113 SOFIA, BULGARIA

FACULTY OF INFORMATION SCIENCES, STATE UNIVERSITY OF LIBRARY STUDIES AND INFORMATION TECHNOLOGIES, SHIPCHENSKI PROHOD 69A, 1574 SOFIA, BULGARIA

E-mail address: `nik@math.bas.bg`